

Theory of Flux-Flow Resistivity near H_{c2} for s -wave Type-II Superconductors

Takafumi Kita

Division of Physics, Hokkaido University, Sapporo 060-0810, Japan

(Dated: February 2, 2008)

This paper presents a microscopic calculation of the flux-flow resistivity ρ_f for s -wave type-II superconductors with arbitrary impurity concentrations near the upper critical field H_{c2} . It is found that, as the mean free path l becomes longer, ρ_f increases gradually from the dirty-limit result of Thompson [Phys. Rev. B1, 327 (1970)] and Takayama and Ebisawa [Prog. Theor. Phys. 44, 1450 (1970)]. The limiting behaviors suggest that $\rho_f(H)$ at low temperatures may change from convex downward to upward as l increases, thus deviating substantially from the linear dependence $\rho_f \propto H/H_{c2}$ predicted by the Bardeen-Stephen theory [Phys. Rev. 140, A1197 (1965)].

PACS numbers: 74.25.Op, 74.25.Fy, 72.10.Bg

Kim *et al.* [1] attributed finite resistivity observed in type-II superconductors to the motion of flux lines, calling it “flux-flow resistivity” ρ_f . Whereas the idea has been accepted widely, we still have poor quantitative understanding of the phenomenon. The early phenomenological theories based on single-flux considerations [2, 3, 4] cannot explain the steep decrease of ρ_f observed near H_{c2} [1]. This region, where microscopic calculations may be performed most easily, has been a subject of later theoretical works [5, 6, 7, 8]. We thereby have a quantitative theory on the flux-flow resistivity near H_{c2} [7, 8], but its validity is still restricted to the dirty limit. Reasonable agreements with the theory have been reported later in a couple of experiments on microwave surface resistivity [9] and flux-flow resistivity [10]. However, the latter experiment found a small discrepancy that ρ_f near T_c is larger than the theoretical prediction [7, 8], which was attributed by Larkin and Ovchinnikov [11] to the extra phonon scattering effective at finite temperatures.

With these backgrounds, this paper provides a quantitative theory on the flux-flow resistivity near H_{c2} applicable to arbitrary impurity concentrations. I thereby hope to establish the domain in which the dirty-limit theory [7, 8] is valid, and look into whether the experiments [9, 10] may be explained by impurity scattering alone. Although a similar study was carried out by Ovchinnikov [12], the behaviors of ρ_f at intermediate impurity concentrations have not been clarified explicitly.

I consider the s -wave pairing with an isotropic Fermi surface and the s -wave impurity scattering in an external magnetic field $\mathbf{H} \parallel \mathbf{z}$. I calculate the complex conductivity $\sigma(\omega)$ below microwave frequencies ω using the quasiclassical equations of superconductivity [13], and finally put $\omega \rightarrow 0$. I use the notation of Ref. 14 but leave the Hall terms out of consideration. Hence I start from the same equations as Eschrig *et al.* [15] in clarifying the motion of a single flux line within the linear-response regime.

The vector potential without perturbation is given by $\mathbf{A}(\mathbf{r}) = Bx\hat{\mathbf{y}} + \tilde{\mathbf{A}}(\mathbf{r})$, where B is the average flux density and $\tilde{\mathbf{A}}$ expresses the spatially varying part of the mag-

netic field satisfying $\int \nabla \times \tilde{\mathbf{A}} d\mathbf{r} = \mathbf{0}$. The corresponding retarded quasiclassical Green’s functions f^R and g^R can be obtained as Ref. 16 with the replacement of the Matsubara frequency ε_n by $-i\varepsilon$. I adopt the units where the energy, length, magnetic field, and electric field are measured by the zero-temperature energy gap $\Delta(0)$ at $H=0$, the coherence length $\xi_0 \equiv \hbar v_F / \Delta(0)$ with v_F the Fermi velocity, $B_0 \equiv \phi_0 / 2\pi \xi_0^2$ with $\phi_0 \equiv hc/2e$ the flux quantum, and $E_0 \equiv \hbar v_F / e \xi_0^2$, respectively. I also put $\hbar = k_B = 1$.

Now, consider the response to a spatially uniform but time-dependent perturbation $\delta \mathbf{A} e^{-i\omega t} = \delta \mathbf{E} e^{-i\omega t} / i\omega$ with $\delta \mathbf{E} \perp \mathbf{H}$. A straightforward calculation based on Eq. (71) of Ref. 14 leads to the following equation for the first-order response $\delta f^R = \delta f^R(\varepsilon, \mathbf{k}, \omega, \mathbf{r})$:

$$\begin{aligned} & \left[-2i\varepsilon + \frac{\langle g_+^R \rangle + \langle g_-^R \rangle}{2\tau} + \hat{\mathbf{v}} \cdot (\nabla - i\mathbf{A}) \right] \delta f^R \\ &= (f_+^R + f_-^R) \hat{\mathbf{v}} \cdot \delta \mathbf{E} / \omega + (g_+^R + g_-^R) \delta \Delta \\ &+ \left(\Delta + \frac{\langle f_+^R \rangle}{2\tau} \right) \delta g^{R\dagger} + \left(\Delta + \frac{\langle f_-^R \rangle}{2\tau} \right) \delta g^R \\ &- \frac{f_+^R \langle \delta g^{R\dagger} \rangle + f_-^R \langle \delta g^R \rangle}{2\tau} + \frac{(g_+^R + g_-^R) \langle \delta f^R \rangle}{2\tau}. \quad (1a) \end{aligned}$$

Here τ is the relaxation time in the Born approximation, $\langle \dots \rangle$ denotes the Fermi-surface average satisfying $\langle 1 \rangle = 1$, and $\Delta(\mathbf{r})$ and $\delta \Delta(\omega, \mathbf{r})$ are the pair potential and its first-order response, respectively. The unit vector $\hat{\mathbf{v}} = \hat{\mathbf{k}}$ specifies a point on the spherical Fermi surface, $f_{\pm}^R \equiv f^R(\varepsilon_{\pm}, \mathbf{k}, \mathbf{r})$ and $g_{\pm}^R \equiv g^R(\varepsilon_{\pm}, \mathbf{k}, \mathbf{r})$ with $\varepsilon_{\pm} \equiv \varepsilon \pm \omega/2$, and the superscript $\dagger$ denotes simultaneous operations of complex conjugation and $(\varepsilon, \mathbf{k}, \omega) \rightarrow (-\varepsilon, -\mathbf{k}, -\omega)$, e.g., $\delta g^{R\dagger}(\varepsilon, \mathbf{k}, \omega, \mathbf{r}) = \delta g^{R*}(-\varepsilon, -\mathbf{k}, -\omega, \mathbf{r})$. Finally, the normalization condition [13] enables us to write δg^R in terms of δf^R as

$$\delta g^R = -(f_+^R \delta f^{R\dagger} + f_-^R \delta f^R) / (g_+^R + g_-^R). \quad (1b)$$

Equation (1) determines the retarded functions δf^R and δg^R for given $\delta \mathbf{E}$ and $\delta \Delta$. In addition, the advanced functions are obtained directly by using Eq. (72) of Ref. 14 as $\delta f^A(\varepsilon, \mathbf{k}, \omega, \mathbf{r}) = \delta f^R(-\varepsilon, -\mathbf{k}, \omega, \mathbf{r})$ and $\delta g^A(\varepsilon, \mathbf{k}, \omega, \mathbf{r}) = -\delta g^{R*}(\varepsilon, \mathbf{k}, -\omega, \mathbf{r})$.

As for the Keldysh functions, I write them following Eschrig *et al.* [15] as $\delta g^K = \delta g^R \phi_- - \phi_+ \delta g^A + \delta g^a$ and $\delta f^K = \delta f^R \phi_- - \phi_+ \delta f^A + \delta f^a$ with $\phi_{\pm} \equiv \tanh(\varepsilon_{\pm}/2T)$. Then a simplified equation results for $\delta g^a = \delta g^a(\varepsilon, \mathbf{k}, \omega, \mathbf{r})$ as

$$\begin{aligned} & \left(-i\omega + \frac{\langle g_+^R \rangle - \langle g_-^A \rangle}{2\tau} + \hat{\mathbf{v}} \cdot \nabla \right) \delta g^a \\ &= (\phi_+ - \phi_-) \left[-(g_+^R - g_-^A) \hat{\mathbf{v}} \cdot \delta \mathbf{E} / \omega + f_+^R \delta \Delta^\dagger - f_-^A \delta \Delta \right] \\ &+ \left(\Delta + \frac{\langle f_+^R \rangle}{2\tau} \right) \delta f^{a\dagger} + \left(\Delta^* + \frac{\langle f_-^A \rangle}{2\tau} \right) \delta f^a \\ &- \frac{f_+^R \langle \delta f^{a\dagger} \rangle + f_-^A \langle \delta f^a \rangle}{2\tau} + \frac{(g_+^R - g_-^A) \langle \delta g^a \rangle}{2\tau}. \end{aligned} \quad (2a)$$

It also follows from the normalization condition that $\delta f^a = \delta f^a(\varepsilon, \mathbf{k}, \omega, \mathbf{r})$ is given in terms of δg^a by

$$\delta f^a = - (f_+^R \delta g^{a\dagger} + f_-^A \delta g^a) / (g_+^R - g_-^A). \quad (2b)$$

Finally, the pair potential $\delta \Delta$ is determined by

$$\delta \Delta = \frac{N(0)V_0}{4i} \int_{-\varepsilon_c}^{\varepsilon_c} d\varepsilon \langle \delta f^R \phi_- - \delta f^A \phi_+ + \delta f^a \rangle, \quad (3)$$

where $N(0)$ is the density of states per spin at the Fermi level, $V_0 > 0$ is the s -wave pairing interaction, and ε_c is the cut-off energy. Once the solution to Eqs. (1)-(3) is obtained self-consistently, the transport current $\delta \mathbf{j} = \delta \mathbf{j}(\omega, \mathbf{r})$ is calculated by

$$\delta \mathbf{j} = -\frac{eN(0)v_F}{2} \int_{-\infty}^{\infty} d\varepsilon \langle \hat{\mathbf{v}} (\delta g^R \phi_- - \delta g^A \phi_+ + \delta g^a) \rangle. \quad (4)$$

The corrections caused by $\delta \mathbf{j}$ to the magnetic field and the charge density may be neglected safely for the relevant weak-coupling superconductors. Equations (1)-(4) are still exact within the linear-response regime and form a basic starting point to calculate complex conductivity of type-II superconductors at arbitrary magnetic fields.

I now concentrate on the region near H_{c2} and solve Eqs. (1)-(3) by expanding every quantity up to second order in $\Delta(\mathbf{r})$ as $f^R = f^{R(1)}$, $g^R = 1 - \frac{1}{2} f^{R(1)} f^{R(1)\dagger}$, $\hat{\mathbf{A}} = \hat{\mathbf{A}}^{(2)}$, $\delta \Delta = \delta \Delta^{(1)}$, $\delta f^{R,a} = \delta f^{R,a(1)}$, and $\delta g^{R,a} = \delta g^{R,a(0)} + \delta g^{R,a(2)}$, with $g^{R(0)} = 0$ as seen from Eq. (1b). The superscript (1) in δf and $\delta \Delta$ will be dropped as it causes no confusions.

The zeroth-order quantity $\delta g^{a(0)}$ is obtained easily from Eq. (2a) as

$$\delta g^{a(0)} = -\frac{2\tau}{1 - i\omega\tau} \frac{\phi_+ - \phi_-}{\omega} \hat{\mathbf{v}} \cdot \delta \mathbf{E}. \quad (5)$$

When put into Eq. (4), this expression yields the normal-state Drude conductivity σ_n , as it should.

Next, Eq. (1a) is simplified for the first-order δf^R into

$$\begin{aligned} & \left[-i\varepsilon + \frac{1}{2\tau} + \frac{\sqrt{B} \sin \theta}{2\sqrt{2}} (e^{-i\varphi} a - e^{i\varphi} a^\dagger) \right] \delta f^R \\ &= \frac{f_+^R + f_-^R}{2} \frac{\hat{\mathbf{v}} \cdot \delta \mathbf{E}}{\omega} + \delta \Delta + \frac{\langle \delta f^R \rangle}{2\tau}, \end{aligned} \quad (6)$$

where $a^\dagger$ and a are creation and annihilation operators satisfying $[a, a^\dagger] = 1$ [16, 17], and (θ, φ) are the polar angles of $\hat{\mathbf{v}}$. Equation (6) can be solved with the Landau-level expansion (LLX) method [17] by expanding $\delta \Delta$ and δf^R in periodic basis functions of the vortex lattice as

$$\delta \Delta(\omega, \mathbf{r}) = \sqrt{V} \sum_{N=0}^{\infty} \delta \Delta_N(\omega) \psi_{N\mathbf{q}}(\mathbf{r}), \quad (7a)$$

$$\delta f^R(\varepsilon, \mathbf{k}, \omega, \mathbf{r}) = \sqrt{V} \sum_{m=-\infty}^{\infty} \sum_{N=0}^{\infty} \delta f_{mN}^R(\varepsilon, \theta, \omega) e^{im\varphi} \psi_{N\mathbf{q}}(\mathbf{r}), \quad (7b)$$

where N denotes the Landau level, $\mathbf{q}$ is an arbitrary chosen magnetic Bloch vector, V is the volume of the system, and $\psi_{N\mathbf{q}}(\mathbf{r})$ satisfies $a\psi_{N\mathbf{q}} = \sqrt{N}\psi_{N-1\mathbf{q}}$ and $a^\dagger\psi_{N\mathbf{q}} = \sqrt{N+1}\psi_{N+1\mathbf{q}}$. The quantities without perturbations are expanded similarly. Near H_{c2} , $\Delta(\mathbf{r})$ may well be approximated using only the $N=0$ level as $\Delta(\mathbf{r}) = \sqrt{V}\Delta_0\psi_{0\mathbf{q}}(\mathbf{r})$. The coefficients Δ_0 and f_{mN}^R have already been obtained in Ref. 16, satisfying $f_{mN}^R = \delta_{mN}\Delta_0\tilde{f}_N^R$ with $\tilde{f}_N^{R*}(\varepsilon, \theta) = \tilde{f}_N^R(-\varepsilon, \theta)$. It then follows from Eqs. (3) and (6) that δf_{mN}^R and $\delta \Delta_N$ may be written as

$$\delta \Delta_N = \Delta_0 \delta_{N1} \delta \tilde{\Delta}_1, \quad (8a)$$

$$\delta f_{mN}^R = \Delta_0 \left(\delta_{m,N-1} \delta \tilde{f}_{N+}^R + \delta_{m,N+1} \delta \tilde{f}_{N-}^R \right). \quad (8b)$$

Equation (6) is thereby transformed into ($\mu = \pm$)

$$\begin{aligned} \sum_{N'} \mathcal{M}_{NN'} \delta \tilde{f}_{N'\mu}^R &= (\tilde{f}_{N+}^R + \tilde{f}_{N-}^R) \delta E_\mu \sin \theta / 4\omega \\ &+ \delta_{N1} \delta_{\mu+} \left(\delta \tilde{\Delta}_1 + \langle \delta \tilde{f}_{1+}^R \rangle / 2\tau \right), \end{aligned} \quad (9)$$

where $E_{\pm} \equiv E_x \pm iE_y$, $\tilde{f}_{N\pm}^R \equiv \tilde{f}_N^R(\varepsilon_{\pm})$, and

$$\mathcal{M}_{NN'} \equiv -i\tilde{\varepsilon} \delta_{NN'} + \beta \sqrt{N+1} \delta_{N,N'-1} - \beta \sqrt{N} \delta_{N,N'+1}, \quad (10a)$$

with $\tilde{\varepsilon} \equiv \varepsilon + i/2\tau$ and $\beta \equiv \sqrt{B} \sin \theta / 2\sqrt{2}$. To solve Eq. (9), let us write the inverse of the matrix $\mathcal{M}$ as

$$K_N^{N'} \equiv (\mathcal{M}^{-1})_{NN'}. \quad (10b)$$

Then $\delta \tilde{f}_{N\mu}^R$ is obtained formally as

$$\delta \tilde{f}_{N\mu}^R = K_N^R \delta E_\mu / \omega + \delta_{\mu+} K_N^1 \left(\delta \tilde{\Delta}_1 + \langle \delta \tilde{f}_{1+}^R \rangle / 2\tau \right), \quad (10c)$$

where K_N^R is defined by

$$K_N^R \equiv \frac{\sin \theta}{4} \sum_{N'} K_N^{N'} (\tilde{f}_{N'+}^R + \tilde{f}_{N'-}^R), \quad (10d)$$

with $\tilde{f}_N^R$ already obtained as [16]

$$f_N^R = K_N^0 / (1 - \langle K_0^0 \rangle / 2\tau). \quad (10e)$$

Taking the angle average of Eq. (10c), we obtain

$$\langle \delta \tilde{f}_{1+}^R \rangle = \frac{\langle K_1^R \rangle}{1 - \langle K_1^1 \rangle / 2\tau} \frac{\delta E_+}{\omega} + \frac{\langle K_1^1 \rangle}{1 - \langle K_1^1 \rangle / 2\tau} \delta \tilde{\Delta}_1. \quad (10f)$$

Equation (10) determines $\delta \tilde{f}_{N\mu}^R$ efficiently and fixes the retarded response δf^R .

We next consider δf^a and substitute Eqs. (5), (7b), and (8b) into Eq. (2b) with $g^R = -g^A = 1$. Expanding δf^a as Eq. (7b), we find that the coefficient δf_{mN}^a can also be written as Eq. (8b). The corresponding $\delta f_{N\mu}^a$ ($\mu = \pm$) is obtained easily as

$$\delta \tilde{f}_{N\mu}^a = K_N^a \frac{\phi_+ - \phi_-}{\omega} \delta E_\mu, \quad (11a)$$

with

$$K_N^a \equiv -\frac{\tau \sin \theta}{2(1 - i\omega\tau)} \left[\tilde{f}_{N+}^R + (-1)^{N-1} \tilde{f}_{N-}^{R*} \right]. \quad (11b)$$

Now we are ready to calculate the pair potential $\delta \Delta$. Let us substitute the above results for $\delta \tilde{f}^R$ and $\delta \tilde{f}^a$ into Eq. (3). Then a straightforward calculation yields the expression for $\delta \tilde{\Delta}_1$ defined by Eqs. (7a) and (8a) as

$$\delta \tilde{\Delta}_1(\omega) = D(\omega) \frac{\delta E_+}{\omega}, \quad (12a)$$

with

$$D(\omega) = \frac{\frac{1}{2i} \int_{-\infty}^{\infty} \left[\frac{\langle K_1^R(\varepsilon, \omega) \rangle}{1 - \langle K_1^1(\varepsilon) \rangle / 2\tau} - \langle K_1^a(\varepsilon, \omega) \rangle \right] \phi(\varepsilon_-) d\varepsilon}{\frac{1}{2} \int_{-\infty}^{\infty} \left\{ \frac{1}{\sqrt{\varepsilon^2 + 1}} + \frac{i \langle K_1^1(\varepsilon) \rangle}{1 - \langle K_1^1(\varepsilon) \rangle / 2\tau} \phi(\varepsilon_-) \right\} d\varepsilon}. \quad (12b)$$

In deriving the formula, use has been made of the symmetries: $K_N^{1*}(\varepsilon) = K_N^1(-\varepsilon)$, $K_N^R(\varepsilon, \omega) = K_N^R(\varepsilon, -\omega) = K_N^{R*}(-\varepsilon, \omega)$, and $K_N^a(\varepsilon, \omega) = (-1)^{N-1} K_N^a(-\varepsilon, \omega) = K_N^{a*}(-\varepsilon, -\omega)$. I also substituted the result $1/N(0)V_0 = \frac{1}{2} \int_{-\varepsilon_c}^{\varepsilon_c} \frac{d\varepsilon}{\sqrt{\varepsilon^2 + 1}}$ at $T = H = 0$ and put $\varepsilon_c \rightarrow \infty$, noting $K_1^1(\varepsilon) \rightarrow i/\varepsilon$ as $\varepsilon \rightarrow \infty$, etc. One can see easily that Eq. (12b) satisfies $D(-\omega) = D^*(\omega)$.

The second-order quantities $\delta g^{R,a(2)}$ can be calculated similarly by expanding them as

$$\delta g^{R,a(2)}(\varepsilon, \mathbf{k}, \omega, \mathbf{r}) = \sum_m \sum_{\mathbf{K}} \delta g_{m\mathbf{K}}^{R,a(2)}(\varepsilon, \theta, \omega) e^{im\varphi + i\mathbf{K} \cdot \mathbf{r}}, \quad (13)$$

where $\mathbf{K}$ is a reciprocal lattice vector of the magnetic Brillouin zone [17]. Since $\delta \mathbf{E}$ is spatially uniform, we only need the $\mathbf{K} = \mathbf{0}$ component within the linear-response regime. It also follows from Eq. (4) that $m \neq \pm 1$ are irrelevant for the current density. Let us substitute the expansions for f^R and δf^R into Eq. (1b) with $g^R = 1$, multiplies it by $e^{\mp i\varphi} / 2\pi V$, and perform integrations over $(\mathbf{r}, \varphi)$. We thereby obtain $g_{\pm}^{R(2)} \equiv g_{m=\pm 1, \mathbf{K}=\mathbf{0}}^{R(2)}$ as

$$\delta g_{\pm}^{R(2)}(\varepsilon, \omega) = \Delta_0^2 \Gamma_{\pm}^R(\varepsilon, \omega) \frac{\delta E_{\mp}}{\omega}, \quad (14a)$$

with ($\mu = \pm$)

$$\Gamma_{\mu}^R \equiv -\frac{1}{2} \sum_N (-1)^N \left[(\tilde{f}_{N+}^R + \tilde{f}_{N-}^R) K_N^R + (\tilde{f}_{N+}^R \delta_{\mu+} + \tilde{f}_{N-}^R \delta_{\mu-}) K_N^1 \frac{\langle K_1^R \rangle / 2\tau + D}{1 - \langle K_1^1 \rangle / 2\tau} \right]. \quad (14b)$$

The quantity $g_{\pm}^{a(2)} \equiv g_{m=\pm 1, \mathbf{K}=\mathbf{0}}^{a(2)}$ can be obtained similarly from Eq. (2a) as

$$\delta g_{\pm}^{a(2)}(\varepsilon, \omega) = \Delta_0^2 \Gamma_{\pm}^a(\varepsilon, \omega) \frac{\phi(\varepsilon_+) - \phi(\varepsilon_-)}{\omega} \delta E_{\mp}, \quad (15a)$$

with

$$\begin{aligned} \Gamma_{\mu}^a &\equiv \frac{\tau \sin \theta}{4(1 - i\omega\tau)} \sum_N (-1)^N [(\tilde{f}_{N+}^R)^2 + (\tilde{f}_{N-}^{R*})^2] \\ &\quad - \frac{\tau \sin \theta}{4(1 - i\omega\tau)^2} \sum_N (-1)^N \langle (\tilde{f}_{N+}^R)^2 + (\tilde{f}_{N-}^R)^2 \rangle \\ &\quad - \frac{\langle \tilde{f}_{0+}^R - \tilde{f}_{0-}^{R*} \rangle K_0^a + (\tilde{f}_{1+}^R \delta_{\mu+} + \tilde{f}_{1-}^{R*} \delta_{\mu-}) \langle K_1^a \rangle}{2(1 - i\omega\tau)} \\ &\quad - \frac{\tau (\tilde{f}_{1+}^R \delta_{\mu+} + \tilde{f}_{1-}^{R*} \delta_{\mu-})}{(1 - i\omega\tau)} D. \end{aligned} \quad (15b)$$

Substituting Eqs. (13)-(15) into Eq. (4) and noting $\Gamma_{\mu}^R(\varepsilon, \omega) = \Gamma_{\mu}^{R*}(-\varepsilon, -\omega)$ and $\Gamma_{\mu}^a(\varepsilon, \omega) = \Gamma_{-\mu}^a(-\varepsilon, \omega) = \Gamma_{\mu}^{a*}(-\varepsilon, -\omega)$, we finally obtain $\delta j_{\pm}^{(2)} \equiv j_{\pm}^{(2)} \pm i j_y^{(2)}$ as

$$\delta j_{\pm}^{(2)}(\omega) = \sigma_f^{(2)}(\omega) \delta E_{\pm}, \quad (16a)$$

with

$$\begin{aligned} \sigma_f^{(2)}(\omega) &\equiv -\Delta_0^2 \frac{eN(0)v_F}{2\omega} \int_{-\infty}^{\infty} d\varepsilon \phi(\varepsilon_-) \\ &\quad \times \sum_{\mu} \langle [\Gamma_{\mu}^R(\varepsilon, \omega) - \Gamma_{\mu}^a(\varepsilon, \omega)] \sin \theta \rangle. \end{aligned} \quad (16b)$$

Thus $\sigma_f^{(2)}$ is isotropic, as expected for any second-order corrections, and satisfies $\sigma_f^{(2)}(-\omega) = \sigma_f^{(2)*}(\omega)$.

Equation (16b) forms the main result of the paper, which is not only exact but also convenient for numerical calculations. The key quantities are $K_N^{N'}$'s defined by Eq. (10b) with Eq. (10a), as may be seen from Eqs. (10d), (10e), (11b), (12b), (14b), and (15b), satisfying $K_N^{N'} = (-1)^{N+N'} K_{N'}^{N'}$. An efficient algorithm to calculate them is obtained as follows. Let us define $\mathcal{D}_N$ ($\tilde{\mathcal{D}}_N$) for $N = 0, 1, 2, \dots$ as the determinant of the submatrix obtained by removing (retaining) the first N rows and columns of the tridiagonal matrix $\mathcal{M}$, Eq. (10a). They satisfy $\mathcal{D}_{N-1} = -i\tilde{\varepsilon} \mathcal{D}_N + \beta^2 N \mathcal{D}_{N+1}$ and $\tilde{\mathcal{D}}_{N+1} = -i\tilde{\varepsilon} \tilde{\mathcal{D}}_N + \beta^2 N \tilde{\mathcal{D}}_{N-1}$ with $\tilde{\mathcal{D}}_1 = -i\tilde{\varepsilon}$ and $\tilde{\mathcal{D}}_0 = 1$, as can be shown with standard matrix manipulations. Then $K_N^{N'}$ for $N' \leq N$ is obtained, by also using standard techniques to invert a matrix, as $K_N^{N'} = \beta^{N-N'} (N!/N')^{1/2} \mathcal{D}_{N+1} \tilde{\mathcal{D}}_{N'}/\mathcal{D}_0$.




FIG. 1: The slope α of the flux-flow resistivity at H_{c2} as a function of the reduced temperature T/T_c for several values of $\xi_E/l \equiv 1/2\pi T_c \tau$ with $\kappa_{GL} = 50$. The curve TTE denotes the prediction of the dirty-limit theory [7, 8].

This algorithm can be put into a more convenient form in terms of $\mathcal{R}_N \equiv \mathcal{D}_{N+1}/\mathcal{D}_N$ and $\tilde{\mathcal{R}}_N \equiv \tilde{\mathcal{D}}_{N-1}/\tilde{\mathcal{D}}_N$ as follows. Let us calculate $\mathcal{R}_N$ and $\tilde{\mathcal{R}}_N$ by

$$\begin{cases} \mathcal{R}_{N-1} = (-i\tilde{\epsilon} + \beta^2 N \mathcal{R}_N)^{-1}, & \mathcal{R}_{N_{\text{cut}}} = i/\tilde{\epsilon}, \\ \tilde{\mathcal{R}}_{N+1} = (-i\tilde{\epsilon} + \beta^2 N \tilde{\mathcal{R}}_N)^{-1}, & \tilde{\mathcal{R}}_1 = i/\tilde{\epsilon}, \end{cases} \quad (17a)$$

for an appropriately chosen large N_{cut} . Then $K_N^{N'}$ for $N' \leq N$ is obtained by

$$\begin{cases} K_0^0 = \mathcal{R}_0, & K_{N'}^{N'} = (\mathcal{R}_{N'}/\tilde{\mathcal{R}}_{N'}) K_{N'-1}^{N'-1}, \\ K_{N+1}^{N'} = \beta \sqrt{N+1} \mathcal{R}_{N+1} K_N^{N'}. \end{cases} \quad (17b)$$

One can check the convergence by increasing N_{cut} . It turns out that $N_{\text{cut}} = 2$ is sufficient both near T_c and in the dirty limit where an analytic calculation is also possible. One can thereby reproduce the formula by Thompson [7] and Takayama and Ebisawa [8] which satisfies $\text{Im} \sigma_f^{(2)}(\omega \rightarrow 0) = 0$. In contrast, $N_{\text{cut}} \gtrsim 1000$ is required in the clean limit at low temperatures.

Using Eq. (16b) and taking the limit $\omega \rightarrow 0$, I have calculated the initial slope of the flux-flow resistivity:

$$\alpha \equiv \left. \frac{H}{\rho_n} \frac{\partial \rho_f}{\partial H} \right|_{H=H_{c2}}, \quad (18)$$

for various impurity concentrations and various values of the Ginzburg-Landau parameter κ_{GL} . The main H dependence in Eq. (16b) lies in $\Delta_0^2 \propto H_{c2} - H$, which can be obtained accurately following Ref. 16.

Figure 1 displays the slope α for several values of $\xi_E/l \equiv 1/2\pi T_c \tau$ with $\kappa_{GL} = 50$. The curve TTE denotes the prediction of the dirty-limit theory by Thompson [7] and Takayama and Ebisawa [8]. Marked mean-free-path dependence is clearly seen. In fact, the slope at $T = T_c$ ($T = 0$) decreases from 4.99 (1.72) in the dirty limit to 0.89 (0.73) at $\xi_E/l = 4.0$. Thus, nonlocal effects in clean systems tend to increase the resistivity

substantially over the prediction of the dirty-limit theory. This result near H_{c2} also suggests that $\rho_f(H)$ at low temperatures may change from convex downward to upward as l increases and may not be fit quantitatively by the Bardeen-Stephen theory $\rho_f \propto H/H_{c2}$ [3]. The slope has also been found to become steeper as κ_{GL} approaches $1/\sqrt{2}$, due to the increase of the coefficient of $\Delta_0^2 \propto H_{c2} - H$, reaching $\alpha = 9.57$ (2.59) at $T = T_c$ ($T = 0$) for $\kappa_{GL} = 1$ and $\xi_E/l = 50$. This fact indicates the necessity of correctly identifying the material parameters, such as κ_{GL} , ξ , and l , for any detailed comparisons between the theory and experiments on the flux-flow resistivity. Besides, a careful experiment will be required, especially near T_c , to determine the slope α which may change appreciably near H_{c2} [9].

In summary, this paper has developed a reliable and efficient method to calculate the flux-flow resistivity near H_{c2} over all impurity concentrations and clarified large dependence of ρ_f on both l and κ_{GL} .

This research is supported by Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

-
- [1] Y. B. Kim, C. F. Hempstead, and A. R. Strnad, Phys. Rev. **139**, A1163 (1965); for a review, see, Y. B. Kim and M. J. Stephen, in *Superconductivity*, edited by R. D. Parks (Dekker, NY, 1969), Vol. 2, p. 1107.
 - [2] M. Tinkham, Phys. Rev. Lett. **13**, 804 (1964).
 - [3] J. Bardeen and M. J. Stephen, Phys. Rev. **140**, A1197 (1965).
 - [4] P. Nozières and W. F. Vinen, Philos. Mag. **14**, 667 (1966).
 - [5] A. Schmid, Phys. Kondens. Materie **5**, 302 (1966).
 - [6] C. Caroli and K. Maki, Phys. Rev. **159**, 306 (1967); **164**, 591 (1967).
 - [7] R. S. Thompson, Phys. Rev. **B1**, 327 (1970).
 - [8] H. Takayama and H. Ebisawa, Prog. Theor. Phys. **44**, 1450 (1970).
 - [9] R. J. Pedersen, Y. B. Kim, and R. S. Thompson, Phys. Rev. **B7**, 982 (1973).
 - [10] R. Meier-Hirmer, M. D. Maloney, and W. Gey, Z. Phys. **B23**, 139 (1976).
 - [11] A. I. Larkin and Y. N. Ovchinnikov, in *Nonequilibrium Superconductivity Vol. 12*, edited by D. N. Langenberg and A. I. Larkin (Elsevier, Amsterdam, 1986) p. 493.
 - [12] Y. N. Ovchinnikov, Zh. Eksp. Teor. Fiz. **66**, 1100 (1974); Sov. Phys. JETP **39**, 538 (1974). His analytic formula for long mean free paths at low temperatures, i.e. Eq. (45), predicts that α should decrease as T increases, which is opposite to what is seen in Fig. 1.
 - [13] For a review, see e.g., J. W. Serene and D. Rainer, Phys. Rep. **101**, 221 (1983).
 - [14] T. Kita, Phys. Rev. **B64** (2001) 054503.
 - [15] M. Eschrig, J. A. Sauls, and D. Rainer, Phys. Rev. **B60**, 10447 (1999).
 - [16] T. Kita, cond-mat/0306231.
 - [17] T. Kita, J. Phys. Soc. Jpn. **67**, 2067 (1998).